

Piedmont Women's Healthcare
A Division of Atlanta Women's Healthcare Specialists
275 Collier Road, N.W., Suite 100-B
Atlanta, GA 30309
404-352-3656 (Phone) | 404-350-5820 (Fax)

www.piedmontwomenshealthcare.com

After Your Endometrial Biopsy: Expectations and Instructions

- 1. **PAIN**: Moderate menstrual-like cramping is normal after having an endometrial biopsy. Take ibuprofen (Motrin) 600mg every 6 hours as needed for this pain. You may also add acetaminophen (Tylenol) 650mg every 6 hours if needed. If your pain is not controlled by these two medications, please call the office.
- 2. **VAGINAL BLEEDING**: You may have menstrual-like vaginal bleeding and pass a few small clots for up to 1 to 2 weeks following your biopsy. However, if you saturate one pad per hour for two hours in a row, or pass clots larger than the size of a tennis ball, please call the office.
- 3. **UNUSUAL DISCHARGE and FEVER**: Please call the office if you have foul-smelling vaginal discharge or a fever of >100.4F or >38C that persists for 4 to 6 hours.
- 4. **ACTIVITY**: You may return to your normal daily activities immediately after your endometrial biopsy.
- 5. **HEALING**: Do not put anything in the vagina for 1 week. In other words, no sex, tampons, or douching.
- 6. **RESULTS**: The results from today's procedure will be available within 7 to 10 business days. Your physician will call you at that time to review the pathology and discuss next steps.

Updated: 04/20/2023